

# Studies on Effect of Different Culture Media on Growth and Phycobilins Content in a Cyanobacterium *Tolypothrix fragilis* (Gardner) Geitler

Janardhan Namdeo Nehul

Department of Botany,

Dada Patil Rajale College of Science, Adinathnagar Tal-Pathardi Dist-Ahmednagar- 414505 (MS), India

**Abstract** - A cyanobacterium *Tolypothrix fragilis* was isolated from the collected soil samples from different locations of Ahmednagar district of Maharashtra state (India). Identification was carried out using morphological variation and taxonomical approaches according to Desikachary. The axenic culture of *Tolypothrix fragilis* was obtained in the laboratory. For the biomass production, different culture media were used namely BG-11, Fogg's medium, Allen and Arnon medium, Zarrouk's medium and CFTRI medium. The biomass was harvested by filtration through double layered muslin cloth and dried using air blower. After harvesting, the biomass obtained was subjected to the growth analysis. Phycobilins were estimated by following the method described by Bennett and Bogorad. Out of the different culture media used, BG-11 medium supported the growth of *Tolypothrix fragilis* properly as compared to other media used. Phycobilins content was found to be more in *Tolypothrix fragilis* grown in Fogg's medium followed by the Zarrouk's medium.

**Key Words**- *Tolypothrix fragilis*, Phycobilins, BG-11, Fogg's medium, Allen and Arnon medium, Zarrouk's medium and CFTRI medium.

## I. INTRODUCTION

Cyanobacteria (blue-green algae, BGA) are morphologically diverse group of phototrophic prokaryotes, which occur in almost every habitat on earth and useful to mankind in various ways [1]. They constitute a vast potential resource in varied applications such as food, feed, fuel, fertilizer, medicine, industry and in combating pollution [1]. Until past few decades of research, cyanobacteria were of academic interests and were mostly ignored as nuisance but, now are proved as potential organisms for much biotechnological utilization [2,3,1]. The interest in these organisms as generators of pharmacologically active and industrially important compounds has been stimulated by recent results [4]. The carbohydrates produced by cyanobacteria have important commercial uses. Since carbohydrates are non-toxic, they are desirable and used in the food industry [5]. Carbohydrates are frequently used in dietary additives for poultry and aquaculture farming [6].

A large number of marine nitrogen-fixing cyanobacteria have been tested for their nutritional value with the hybrid *Tilapia* fish fry [7]. The marine cyanobacterium *Phormidium valderianum* BDU 30501 has shown to serve as a complete aquaculture feed source, based on the nutritional qualities and non-toxic nature with animal model experiments [1]. Several micro algae such as *Chlorella*, *Scenedesmus* and *Coelastrum* have been established as good quality protein sources [8]. The main advantage of these species is their high protein content, therefore they are used as food supplements. They also present great benefits to human health due to their antioxidant properties, their role as activator of cell regeneration, and their positive effect on kidney and memory problems [9].

Cyanobacteria possess all the known phycobiliproteins such as phycocyanin, phycoerythrin, Phycoerythrocyanin and allo-phycocyanin. Among them, Phycocyanin and phycoerythrin are commercially valuable. Linablue, a phycocyanin product from Dainippon Ink and chemicals Inc., Japan is an odorless, non-toxic blue powder and used for coloring candy, ice-cream, dairy products and soft drinks [10]. Phycoerythrin from *Spirulina* and other cyanobacteria is used as a food colour for products like ice-cream. [11].

Apart from the use of phycobilins as food grade dyes, they are also used as tools for basic research and medical diagnostics. They are used in fluorescence microscopy and fluorescence immunoassays

[12,13].Phycocyanin, the major phycobiliprotein also exhibited anti-cancer activity, stimulation of immune system and ability to treat ulcers and haemorrhoidal bleeding [2].The present study was carried out for the estimation of growth and phycobilins content in *Tolypothrixfragilis* in different culture media.

## II. MATERIALS AND METHOD

*A. Method of collection*-The soil samples from 5-10 cm deep soil layers were collected using the scalpels. Soil samples were collected in polythene bags of size 6 x 4 inches.

*B. media*-The different culture media namely BG-11[14]; Fogg's medium [15,16]; Allen and Arnon's medium [17; CFTRI medium [18]) and Zarrouk's medium [19] were used for the rich growth of *Tolypothrixfragilis*. These media were separately used in different sets.

*C. Isolation of cyanobacterial species*-The dry soil samples were spread in petri dishes and moistened with sterilized distilled water and cultures were incubated in light. When the visible growth of cyanobacteria begins to appear in the cultures, these cultures were used for the isolation of unialgal cultures of *Tolypothrixfragilis*.

*D. Identification of the algal samples*-Morphometric studies were carried out by using ocular and stage micrometer. The identification of *Tolypothrixfragilis* was carried out using monograph and keys of Desikachary[20].

*E. Biomass production*-For production of biomass, glass bottles (300 mL capacity) were used. The bottles were filled with 100 mL medium and autoclaved. The inoculum was ground in the sterile mortar and pestle in laminar air flow. Then the bottles were inoculated with 5 mL of unialgal suspension of *Tolypothrixfragilis* and labeled properly. All the cultures were maintained in the culture room at temperature  $28 \pm 2^\circ\text{C}$  under 8-h light/16-h dark photoperiod with a photosynthetic photon flux density of  $40 \mu\text{moles}^{-2}\text{S}^{-1}$  provided by cool white fluorescent tube lights. After harvesting, the biomass obtained was subjected to the growth and proteins analysis.

### *F. Estimation of phycobilins*

Phycobilins were estimated by following the method described by Bennett and Bogorad[21]. Two mL cell suspension was centrifuged at 4500 rpm for 10 minutes and pellet of algal biomass was used for the estimation of total phycobilins. The pellet of biomass of *Tolypothrixfragilis* was blotted with filter papers to remove maximum quantity of water from it. The pellet thus dried was homogenized in 100  $\mu\text{l}$  glycerol and placed in the dark for 3 hours. After the contents were mixed, 5 mL of 10 % ammonium sulphate containing 1mL of 3 mM sodium azide and 1mL of 10 mM  $\text{Na}_2\text{EDTA}$  [22]. The resultant cell suspension was sonicated for 10 minutes and then centrifuged at 4500 rpm for 10 minutes. Absorbance of the supernatant solution was determined at 665 nm, 620 nm and 650 nm on UV Visible spectrophotometer (Systronics, India; model 2202). The amounts of phycoerythrin, phycocyanin and allophycocyanin were calculated by using following equations [21].

A. Amount of Phycoerythrin (PE)

$$= A_{565} \cdot 2.8 [\text{PC}] - 1.34 [\text{APC}] \text{ mg ml}^{-1} / 12.7$$

B. Amount of Phycocyanin (PC),

$$= A_{620} - 0.7 \times A_{650} \text{ mg ml}^{-1} / 7.38$$

C. Amount of allophycocyanin (APC)

$$= A_{650} \cdot 0.19 \times A_{620} \text{ mg ml}^{-1} / 5.65$$

Where,  $A_{565}$  = Absorbance of supernatant solution at 565nm,

$A_{620}$  = Absorbance of supernatant solution at 620 nm,

$A_{650}$  = Absorbance of supernatant solution at 650nm.

D.Sum of PE, PC and APC gives content of total phycobiliproteins.

Amount of Phycobiliproteins are expressed as % of dry weight basis.

### III. RESULTS AND DISCUSSION

Out of the different culture media used, BG-11 medium supported the growth of *Tolypothrixfragilis* properly as compared to other media used. Allen and Arnon medium also supported growth but after 20 to 25 days, photo bleaching of biomass was observed. Other growth media, such as Fogg's medium and Zarrouk's medium supported the growth of *Tolypothrixfragilis* but the growth rate was very slow.

Yield of biomass is one of the direct measures of quantity of biomass produced per unit area within a specific time. Higher yield indicates higher biomass produced per unit area. Comparison of *Tolypothrixfragilis* in different media showed that highest biomass per bottle in terms of dry weight was produced in BG-11 medium followed by Allen and Arnon medium. The phycobilins content was found to be more in the *Tolypothrixfragilis* grown in Fogg's medium followed by the Zarrouk's medium. CFTRI medium showed poor response for the phycobilins content.

Table-1. Influence of different media on growth and phycobilins in *Tolypothrixfragilis*

Sr.no	Medium	Fresh wt. (g)	Dry wt. (g)	Phycobilins %
1	BG-11	1.98±0.17 <sup>a</sup>	0.18±0.06 <sup>a</sup>	3.76±0.03 <sup>c</sup>
2	Allen & Arnon	1.74±0.19 <sup>a</sup>	0.17±0.04 <sup>a</sup>	3.83±0.02 <sup>c</sup>
3	Fogg's Medium	1.22±0.25 <sup>c</sup>	0.10±0.03 <sup>c</sup>	5.37±0.04 <sup>a</sup>
4	Zarrouk' Medium	1.21±0.12 <sup>c</sup>	0.10±0.04 <sup>c</sup>	4.17±0.03 <sup>b</sup>
5	CFTRI	1.37±0.19 <sup>b</sup>	0.12±0.04 <sup>b</sup>	2.88±0.02 <sup>d</sup>

Values are mean±SE of three independent experiments.

Cyanobacteria are photoautotrophic bacteria and require all the essential major and minor elements. The heterocystous cyanobacteria fix atmospheric nitrogen and they can use atmospheric nitrogen as a source of nitrogen. In bottles, the medium does not come in contact with atmospheric nitrogen and the source needs to be added in the culture medium. If the culture medium is devoid of nitrogen, it results in poor growth of cyanobacteria. Similar results were reported by Olatz[23]; medium lacking nitrogen source, results in yellowish green color of the cells which is a characteristic of nitrogen deficiency. In the culture methods like photo- bioreactors, pure nitrogen is continuously bubbled into culture medium, [24,25,26]. so that cultures do not get affected due to nitrogen deficiency.

The growth of *Tolypothrixfragilis* was more in BG-11 medium than in other media. For optimum growth of cyanobacteria, appropriate  $Ka^+ : Na^+$  ratio is required in the cytoplasm. Adequate  $Na^+$  is required by nitrogen fixing cyanobacteria for conversion of molecular nitrogen into ammonia [27]. BG-11 medium consists moderate concentration of  $Na^+$  and in Allen and Arnon medium, Zarrouk's medium and CFTRI medium there is high concentration of  $Na^+$  while in Fogg's medium; there is no  $Na^+$  source. *Tolypothrixfragilis* from moist soil habitat, which may not require high concentration of  $Na^+$  ions in the medium.

Composition and pH of the medium also affect the content of phycobiliprotein and there is change in the colour of culture of cyanobacteria. The production of phycobilins is greatly affected by the pH. This indicates the inability of the cyanobacteria to maintain a constant internal pH [28].

## IV.CONCLUSION

There is an effect of different culture media on growth. It may be because of either composition of culture medium or pH value. The percentage of phycobilins also changed in various culture media used.

## REFERENCES

- [1] Thajuddin N, Subramanian G, "Cyanobacterial biodiversity and potential applications in biotechnology", Curr. Sci. 89: 47-57,2005.
- [2] Richmond A. "In: *Algal Biotechnology*", (Eds.). Round FE, Chapman DJ, Bristol UK: Biopress Ltd. pp. 269-330,1990.
- [3] Sundararaman M, Sekar S, "Biotechnological potential of cyanobacteria." In: *Algal biotechnology*, Trivedi PC (ed.), Pointer Publishers, Jaipur, India. pp 31–52,2001.
- [4] Singh DP, Tyagi MB, Kumar A, Prasuna EG, "Bioactive Secondary Metabolites from cyanobacteria." Pg.275-292. In *Algal research in India*. Edt. N. Anand, Pbl. Bsmp's Dehradun. India,2002.
- [5] Bauernfeind JC, "Carotenoids as colorants and Vitamins A Precursors". London: Academic Press,1981.
- [6] Hirschberg J, Chamoritz D (1994) In: *The Molecular Biology of Cyanobacteria*, (Ed.). Bryant, DADordrecht, The Netherlands: Kluwer Academic Publishers. pp.559-579, 1994.
- [7] Mitsui A, Entermann B, Gill K, "Indoor and outdoor cultivation of *Tilapia* in seawater with algae as a sole food source". In *Proceedings of the 2nd North Pacific Aquaculture System*, Tokyo University, Japan, pp. 323-340
- [8] Anusuya. DM, Subbulaxmi G, Madhavi DM, Venkataraman LV, "Studies on the proteins of mass cultivated blue green alga (*Spirulina platensis*)". J. Agric. Food Chem. 29:522-525,1981.
- [9] Gutiérrez-Rebolledo, G.A.; Galar-Martínez, M.; García-Rodríguez, R.V.; Chamorro-Cevallos, G.A.; Hernández-Reyes, A.G.; Martínez-Galero, E., "Antioxidant effect of *Spirulina (Arthrospira) maxima* on chronic inflammation induced by Freund's complete adjuvant in rats". J. Med. Food, 18, 865–871,2015.
- [10] Cohen Z, "Products from microalgae". In *handbook of microalgal Mass Culture*". (Eds.). Richmond, A. Boca Raton FL, USA: CRC Press, pp. 421-454,1986.
- [11] Borowitzka MA, "In: Proceedings of the first Asia-Pacific Conference on Algal Biotechnology". (Eds.) Phang SM, Lee YK, Borowitzka MA and Whitton BA, University of Malaya, Kuala Lumpur, pp. 5-15,1994.
- [12] Glazer AN, Stryer L, "Phycofluor probes" Trends Biochem. Sci. 9:423-427,1984
- [13] Kronick MN, "The use of phycobiliproteins as fluorescent labels in immunoassay". J Immunol Methods 92:1-13, 1986.
- [14] Rippka R, Deruelles J, Waterbury JB, Herd man M, Stainer RY, "Generic assignments. Strain histories and properties of pure cultures of cyanobacteria." J. Gen. Microbiol. 111: 1-61,1979.
- [15] Fogg GE, "Growth and heterocyst production in *Anabaena cylindrica* Lemm. II. In relation to carbon and nitrogen metabolism". Ann. Bot. 13: 241-259,1949.
- [16] Jacobson L, "Maintenance of iron supply in nutrient solutions by a single addition of ferric potassium ethylene diamine tetra-acetate". Plant Physiology 26: 411-413,1951.
- [17] Allen MB, Arnon DI, "Studies on nitrogen fixing blue green algae. I. Growth and nitrogen fixation in *Anabaena cylindrica*." Plant Physiol. 30:366-372, 1955.
- [18] Venkatarama, LV, Becker EW, "Production and utilization of the blue-green alga *Spirulina* in India". Biomass4: 105-125,1984.
- [19] Zarrouk C, "Contribution a l'etude d'uncyanophyce. Influence de divers' facteurs physiques et chimiques sur la croissance et la photosyntheses de *Spirulinamaxima*". Ph.D. Thesis. University of Paris, France,1966.
- [20] Desikachary TV, "A monograph on Cyanophyta", ICAR Publication, New Delhi. 1-686,1959.
- [21] Bennett A, Bogorad L, "complementary chromatic adaptation in a filamentous blue-green alga". The Journal of Cell Biology 58: 419-435,1973.
- [22] Whyman M, "An *in vitro* method for estimation of phycoerythrin concentration in marine cyanobacteria (*Syneccoccus* Sp.)", Limol. Oceanogr 37: 1300-1306, 1992.
- [23] Olatz F, Ramon G, Juan LS, "Carotenoides composition in the Cyanobacterium *Phormidium laminosum*. Effect of nitrogen starvation". FEBS Letters, 282: 300-304,1991.
- [24] Humberto J, Silva IC, Rodolfo JE, "Effect of nutritional factors on the culture of *Nostoc* sp. as a source of phycobiliproteins." Appl Microbiol Biotechnol 31: 293-297,1989.
- [25] Vonshak A, "*Spirulina platensis* (Arthrospira): Physiology, Cell Biology and Biotechnology." Taylor & Francis, London. ISBN 0-7484-0674-3,1993.
- [26] Roxana Olvera-Ramirez, Marisela Coria-Cedillo, Rosa Olivia Canizares-Villanueva Fernando Martinez-Jeronimo, Teresa Ponce-Noyola, Elivira Rios-Leal, "Growth evaluation and byproduct characterization of *Calothrix* sp.". Bioresource Technology 72: 121-124,2000.
- [27] Becker EW, "In: Microalgae - Biotechnology and Microbiology", Cambridge, Cambridge University Press, ISBN 0- 521 35020-4, pp 1-4,1994.
- [28] Roe, S., "Protein Purification Techniques: A practical Approach". Oxford University press, Oxford, UK.,2000.